



**Europäisches  
Patentamt**


**European  
Patent Office**

**PCT/NL**

**Office européen  
des brevets**

**10/528545**  
**03/00661**

**06.11.03**



**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**

**02078992.1**

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**



Anmeldung Nr:  
Application no.: 02078992.1  
Demande no:

Anmeldetag:  
Date of filing: 26.09.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Stichting  
Researchfonds Pathologie  
Jekerstraat 19 2HG  
1078 LW Amsterdam  
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Method for detecting and typing of cutaneous HPV and primers and probes for use therein

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C12Q1/70

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

26. 09. 2002

P60890EP00

(92)

Title: Method for detecting and typing of cutaneous HPV and primers and probes for use therein.

### Field of the invention

The invention relates to a method for detecting and typing of human papillomavirus (HPV). In particular, the invention relates to a method for detecting and typing cutaneous HPVs of supergroup B. Also, the invention  
5 relates to primers and probes for use in a method of the invention.

### Background of the invention

The etiologic agents of all warts are a diverse group of viruses known as human papillomaviruses (HPVs) that, in all, constitute a group of more  
10 than 100 types of viruses, as identified by variations in DNA sequence. The various HPVs cause a variety of cutaneous and mucosal diseases. Certain types may cause warts, or papillomas, which are benign (noncancerous) tumors. Others have been found to cause invasive carcinoma of the uterine cervix.

15 Of all HPV types, more than 30 can be passed from one person to another through sexual contact. HPV infection is thereby one of the most common sexually transmitted diseases (STDs). Lesions may appear within several weeks to several months after sexual contact with a person who has HPV, or they may never appear, since HPV infections often do not cause any  
20 symptoms. There is currently no medical cure to eliminate a papillomavirus infection, although the lesions and warts themselves can be treated.

Papillomaviruses together with the Polyomaviruses belong to the Papovaviridae, a family of small, non-enveloped tumor viruses with a double stranded DNA genome. Individual isolates are highly host specific.  
25 Papillomaviruses specific for human (HPV), bovine (BPV), avian (e.g. FPV),

deer (DPV), Cottontail rabbit (CRPV), European elk (EEPV) and numerous other animals have been identified.

Unlike other viruses that are mostly grouped by serotypes, papillomaviruses are classified by genotypes for the same host and are  
 5 numbered in the order of identification (e.g. HPV1-HPV83 and BPV1-BPV4).

HPVs are broadly classified into low-risk and high-risk types, based on their ability to induce malignant changes in infected cells. Low risk HPV types such as 1, 2, 4, 6, 11, 13 and 32 are primarily associated with benign lesions or common warts while the high risk types, such as 16, 18, 31, 33, 35,  
 10 39, 45, 51, 52, 56, 58, 59, 68, and 69 are primarily associated with premalignant and malignant epithelial lesions. These high-risk types of HPV cause growths that are usually flat and nearly invisible, as compared with the warts caused by low-risk types, e.g. HPV-6 and HPV-11.

Papillomaviruses can in general also be classified according to tissue  
 15 tropism. Although all known human papillomaviruses strictly infect cells of epithelial origin (keratinocytes) and induce 'epithelial tumors' they can be grouped according to their tissue tropism into either mucosal or cutaneous HPV types.

The mucosal types (e.g. HPV6 and HPV16), include those HPVs that  
 20 infect mucosa of the genital and respiratory tracts. The cutaneous types (e.g. HPV1 and HPV8), include those HPVs that infect generally the external skin, including those that cause cutaneous warts and those that cause skin carcinomas in patients suffering from epidermodysplasia verruciformis (EV; a rare hereditary disease characterized by disseminated flat warts and macular  
 25 lesions arising during childhood and by a predisposition to HPV-induced skin carcinoma later in life).

The HPV types that have been shown to be associated with EV constitute a group of cutaneous HPVs classified as EV-type (e.g. types 5, 8, 17,  
 20 and 36). Some of these viruses may cause cancer in genetically predisposed  
 30 individuals only. Based on their phylogenetic relationship the EV-type HPVs

together with the EV-type related HPVs and type 4 HPVs are collectively classified as the supergroup B-type and form the largest group of cutaneous HPV types.

5 The above is articulated to indicate that, generally, the type of HPV predicts the site of the infection and its clinical course.

Recent data suggest that the supergroup B-type cutaneous human papillomaviruses might be involved in the pathogenesis of non-melanoma skin cancer (NMSC), particularly in immunosuppressed (e.g. renal transplant recipients) patients. In addition, cutaneous HPV DNA, especially of types 5, 36  
10 and 38 can be detected in up to 90% of psoriasis lesions, suggesting an etiological relationship. Studies on cutaneous HPVs are, however, hampered by the fact that these comprise a large and highly heterogeneous virus group for which a plurality of PCR-based detection systems in parallel need to be employed.

15 The cutaneous HPV detection and typing systems of the prior art have a number of limitations. One disadvantage of the HPV detection and typing systems of the prior art is that only a restricted number of HPV types can be detected in a single assay. Therefore several separate PCR assays need to be performed to cover a broad range of different types (see e.g. Shamanin *et al.*, 1996, J. Natl. Cancer Inst. 88: 802-811; Meijer *et al.*, 2001, Cancer Detect.  
20 Prev. 25: 533-547).

Further, the assays of the prior art require relatively large DNA fragments (Shamanin *et al.*, 1996, J. Natl. Cancer Inst. 88: 802-811; Berkhout *et al.*, 1995, J. Clin. Microbiol. 33: 690-695; Forslund *et al.*, 1999, J. Gen. Virol.  
25 80: 2437-2443; Meijer *et al.*, 2001, Cancer Detect. Prev. 25: 533-547). Since both appropriate preservation of material (e.g. formalin-fixation) as well as poor preservation of clinical material may yield degraded DNA, such assays can only suitably be performed on relatively fresh, unfixed material. This hampers their use and prevents standardization of methodologies.

Moreover, the current assay systems that are suitable for detection of cutaneous HPVs have no system for typing coupled thereto. As a result, laborious reamplification, cloning and sequencing of DNA amplification products is required for typing of HPVs (Shamani *et al.*, 1996, J. Natl. Cancer  
5 Inst. 88: 802-811; Berkhout *et al.*, 1995, J. Clin. Microbiol. 33: 690-695).

Another problem of some methods for detection of multiple HPV types at respectable sensitivity of the prior art is that their format is a so-called nested-PCR format (Berkhout *et al.*, 1995, J. Clin. Microbiol. 33: 690-695). Such methods are extremely sensitive to contamination, as very small  
10 amounts of contaminating template DNA will already yield false positive results. Moreover, such a format is very laborious.

#### Summary of the invention

In order to overcome these difficulties the inventors have now found a  
15 method for the detection of any of the various cutaneous HPV types that belong to supergroup B. Furthermore, and in an additional aspect of the invention, the inventors have found a method for an efficient typing system for HPV types belonging to supergroup B that is compatible with the method for detection.

20 A method for the detection of cutaneous supergroup B HPVs according to the present invention comprises the steps of providing a sample suspected of harbouring cutaneous supergroup B HPVs, providing a plurality of pairs of bi-directional primers collectively substantially complementary to DNA of all cutaneous supergroup B HPVs, performing a reaction to amplify  
25 DNA derived from the said sample using said plurality of primers; and detecting DNA amplification products from cutaneous supergroup B HPV by hybridising the reaction products of the said DNA amplification reaction to a plurality of generic cutaneous supergroup B HPV probes.

In an additional aspect, the invention provides a method for typing of  
30 a cutaneous supergroup B HPV comprising the steps of providing DNA

amplification products by amplifying DNA of cutaneous supergroup B HPV between a plurality of pairs of bi-directional primers, and detecting DNA amplification products from one or more supergroup B HPV types by hybridising the said amplification products to at least one cutaneous  
5 supergroup B HPV probe that is substantially complementary to the DNA of at least one but not all cutaneous supergroup B HPV types.

In a further aspect, the invention provides a set of bi-directional amplification primers for the amplification of supergroup B HPV-DNA.

In another aspect, the invention provides generic oligonucleotide  
10 probes for the generic detection of supergroup B HPV-DNA.

In yet another aspect, the invention provides type-specific oligonucleotide probes for the type-specific or selective detection of individual HPV-types belonging to supergroup B of HPV.

15 Description of the drawings

Figure 1 presents suitable pairs of bi-directional primers for the amplification of DNA from cutaneous HPV belonging to supergroup B.

Figure 2 presents the sequences of supergroup B HPV-specific primer-binding regions in DNA of cutaneous HPV belonging to supergroup B (5'- 3').

20 For clarity reasons and to facilitate alignment of the sequences presented, the complementary strands of these regions are not shown, but are understood to be included.

Figure 3 presents suitable generic detection probes for detection of amplicons derived from cutaneous supergroup B HPV (5'DIG-3'). As illustrated  
25 in the text, these probes may also consist of the complementary strands of those shown.

Figure 4 presents the sequences of probe-binding region in DNA of cutaneous supergroup B HPV (5'- 3'). For clarity reasons and to facilitate alignment of the sequences presented, the complementary strands of these  
30 regions are not shown, but are understood to be included.

Figure 5 presents type-specific detection probes suitable for reverse line blot (RLB) typing of amplicons derived from cutaneous supergroup B HPV (5' amino-3'). As illustrated in the text, these probes may also consist of the complementary strands of those shown.

5           Figure 6 presents the double stranded DNA sequence of HPV type 4 (numbering according to EMBL accession number X70827). Indicated are positions of bi-directional primers (underlined) and probe-binding region (shaded) as defined in Figures 2 and 4, respectively.

10   Detailed description of the invention

A "complement" or "complementary sequence" is a sequence of nucleotides which forms a hydrogen-bonded duplex with another sequence of nucleotides according to Watson-Crick base-pairing rules. For example, the complementary base sequence for 5'-AAGGCT-3' is 3'-TTCCGA-5'.

15           "Expression" refers to the transcription of a gene into structural RNA (rRNA, tRNA) or messenger RNA (mRNA) and, if applicable, subsequent translation into a protein.

Polynucleotides are "heterologous" to one another if they do not naturally occur together in the same organism. A polynucleotide is  
20   heterologous to an organism if it does not naturally occur in its particular form and arrangement in that organism.

Polynucleotides have "homologous" sequences if the sequence of nucleotides in the two sequences is the same when aligned for maximum correspondence as described herein. Sequence comparison between two or  
25   more polynucleotides is generally performed by comparing portions of the two sequences over a comparison window to identify and compare local regions of sequence similarity. The comparison window is generally from about 20 to 200 contiguous nucleotides. The "percentage of sequence homology" for  
polynucleotides, such as 50, 60, 70, 80, 90, 95, 98, 99 or 100 percent sequence  
30   homology may be determined by comparing two optimally aligned sequences



over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may include additions or deletions (i.e. gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is  
5 calculated by: (a) determining the number of positions at which the identical nucleic acid base occurs in both sequences to yield the number of matched positions; (b) dividing the number of matched positions by the total number of positions in the window of comparison; and (c) multiplying the result by 100 to yield the percentage of sequence homology. Optimal alignment of sequences for  
10 comparison may be conducted by computerized implementations of known algorithms, or by inspection. Readily available sequence comparison and multiple sequence alignment algorithms are, respectively, the Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. 1990. J. Mol. Biol. 215:403; Altschul, S.F. et al. 1997. Nucleic Acid Res. 25:3389-3402) and  
15 ClustalW programs both available on the internet. Other suitable programs include GAP, BESTFIT and FASTA in the Wisconsin Genetics Software Package (Genetics Computer Group (GCG), Madison, WI, USA).

As used herein, "substantially complementary" means that two nucleic acid sequences have at least about 65%, preferably about 70%, more  
20 preferably about 80%, even more preferably 90%, and most preferably about 98%, sequence complementarity to each other. This means that the primers and probes must exhibit sufficient complementarity to their template and target nucleic acid, respectively, to hybridise under stringent conditions. Therefore, the primer sequences as disclosed in this specification need not  
25 reflect the exact sequence of the binding region on the template and degenerate primers can be used. A substantially complementary primer sequence is one that has sufficient sequence complementarity to the amplification template to result in primer binding and second-strand synthesis.

The term "hybrid" refers to a double-stranded nucleic acid molecule, or duplex, formed by hydrogen bonding between complementary nucleotides. The terms "hybridise" or "anneal" refer to the process by which single strands of nucleic acid sequences form double-helical segments through hydrogen  
5 bonding between complementary nucleotides.

The term "oligonucleotide" refers to a short sequence of nucleotide monomers (usually 6 to 100 nucleotides) joined by phosphorous linkages (e.g., phosphodiester, alkyl and aryl-phosphate, phosphorothioate), or non-phosphorous linkages (e.g., peptide, sulfamate and others). An oligonucleotide  
10 may contain modified nucleotides having modified bases (e.g., 5-methyl cytosine) and modified sugar groups (e.g., 2'-O-methyl ribosyl, 2'-O-methoxyethyl ribosyl, 2'-fluoro ribosyl, 2'-amino ribosyl, and the like). Oligonucleotides may be naturally-occurring or synthetic molecules of double- and single-stranded DNA and double- and single-stranded RNA with circular,  
15 branched or linear shapes and optionally including domains capable of forming stable secondary structures (e.g., stem-and-loop and loop-stem-loop structures).

The term "primer" as used herein refers to an oligonucleotide which is capable of annealing to the amplification target allowing a DNA polymerase to  
20 attach thereby serving as a point of initiation of DNA synthesis when placed under conditions in which synthesis of primer extension product which is complementary to a nucleic acid strand is induced, i.e., in the presence of nucleotides and an agent for polymerization such as DNA polymerase and at a suitable temperature and pH. The (amplification) primer is preferably single  
25 stranded for maximum efficiency in amplification. Preferably, the primer is an oligodeoxy ribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact lengths of the primers will depend on many factors, including temperature and source of primer. A "pair of bi-directional primers" as used

herein refers to one forward and one reverse primer as commonly used in the art of DNA amplification such as in PCR amplification.

The term "probe" refers to a single-stranded oligonucleotide sequence that will recognize and form a hydrogen-bonded duplex with a complementary  
5 sequence in a target nucleic acid sequence analyte or its cDNA derivative.

The terms "stringency" or "stringent hybridization conditions" refer to hybridization conditions that affect the stability of hybrids, e.g., temperature, salt concentration, pH, formamide concentration and the like. These conditions are empirically optimised to maximize specific binding and minimize non-  
10 specific binding of primer or probe to its target nucleic acid sequence. The terms as used include reference to conditions under which a probe or primer will hybridise to its target sequence, to a detectably greater degree than other sequences (e.g. at least 2-fold over background). Stringent conditions are sequence dependent and will be different in different circumstances. Longer  
15 sequences hybridise specifically at higher temperatures. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridises to a perfectly matched probe or  
20 primer. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M  $\text{Na}^+$  ion, typically about 0.01 to 1.0 M  $\text{Na}^+$  ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes or primers (e.g. 10 to 50 nucleotides) and at least about 60°C for long probes or primers (e.g. greater than 50  
25 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringent conditions or "conditions of reduced stringency" include hybridization with a buffer solution of 30% formamide, 1 M NaCl, 1% SDS at 37°C and a wash in 2x SSC at 40°C. Exemplary high stringency conditions include hybridization in 50%  
30 formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1x SSC at 60°C.

Hybridization procedures are well known in the art and are described in e.g. Ausubel et al, Current Protocols in Molecular Biology, John Wiley & Sons Inc., 1994.

5 A method of the present invention permits the detection, identification and diagnosis of human papillomaviruses belonging to supergroup B and human papillomavirus infections in organisms susceptible to such infections that are caused by supergroup B HPV.

#### Detection of supergroup B HPVs

10 A method for detection of supergroup B HPVs according to the present invention may be performed on samples derived from humans, but also on other samples, such as environmental samples, or samples of other localities where HPV may reside, such as samples from other organisms such as insects. Preferably, a sample for use in a method of the invention comprises  
15 a human sample, such as a sample of a patient's cells. More preferably, the sample comprises the patient's cells from the site of a potential or suspected infection.

The sample of a patient's cells may comprise a sample of tumorous skin cells (e.g. in the case of warts, verrucas and the like, presumably caused  
20 by cutaneous HPV infections). Alternatively, the sample may comprise a sample of tumorous mucosal cells, such as cervical cells in the case of HPV infections of the female urinogenital tract, but also cancerous tissue of lung, larynx, oral cavity, tonsil, oesophagus, penis, vulva, anus, rectum, prostate, ovary and other tissues may be used as a sample in a method of the present  
25 invention.

Methods of obtaining and preparing such samples for use in the method of the invention are known to those skilled in the art or will be apparent from the present disclosure. For general methods concerning DNA analysis and manipulations see e.g. Molecular Cloning: A Laboratory Manual,  
30 2nd Ed., Vol. 1-3, eds. Sambrook et al. Cold Spring Harbor Laboratory Press

(1989) or Current Protocols in Molecular Biology, eds. Ausubel et al., Greene Publishing and Wiley-Interscience, New York (1987) and periodic updates thereof.

5 Preferably, a method according to the invention is used for the detection of HPV types 4, 5, 8, 9, 12, 14, 15, 17, 19, 20, 21, 22, 23, 24, 25, 36, 37, 38, 47, 48, 49, 50, 60 and/or 65. All these types are classified as cutaneous supergroup B HPV.

10 It is an advantageous aspect of the invention that members of the cutaneous HPV supergroup B that are not yet discovered may suitably be detected by using a method of the present invention. Moreover, the method of typing of supergroup B HPVs may be expanded to facilitate for the incorporation of these newly discovered types in a HPV typing system according to the present invention.

15 The inventors have now found two discrete regions in the DNA of all cutaneous supergroup B HPV types that are closely spaced together and that appear homologous between the various supergroup B HPV types. These two homologous regions, or consensus regions, are located in the L1 open reading frame (ORF) of the cutaneous supergroup B HPV genome. The DNA sequence of these two consensus regions is presented in Figure 2. The first consensus  
20 region comprises a region defined herein as the region extending from about nucleotide position 6539 to about nucleotide position 6559 in a 5'-3' direction on the genome of HPV type 4 (numbering according to EMBL accession number X70827, HPV type 4) and the corresponding region within the genome of other cutaneous supergroup B HPV types. The second consensus region  
25 comprises a region defined herein as the region extending from about nucleotide position 6583 until about nucleotide position 6610 in a 5'-3' direction on the HPV 4 genome and the corresponding region within the genome of other cutaneous supergroup B HPV types.

30 Based on this finding the present invention provides a set of bi-directional primers wherein each individual primer is capable of annealing to

one of the two consensus regions of a plurality of cutaneous supergroup B HPV types. A complete set preferably provides a plurality of pairs of bi-directional primers that, when used in concert, are capable of annealing to these two consensus regions in the DNA of any cutaneous supergroup B HPV and of  
5 amplifying the defined DNA sequences between and adjacent these two consensus regions, i.e. flanked by these two consensus regions.

An example of suitable pairs of bi-directional primers to amplify the region between the above mentioned two consensus regions from all cutaneous supergroup B HPVs is presented in Figure 1. From the present disclosure it  
10 will be apparent to the person skilled in the art that other sets of primers than those of Figure 1 may be designed and used for amplification of specific sequences flanked by or flanking the two consensus regions in the L1 ORF of the cutaneous supergroup B HPV genome. For instance, it is possible to use somewhat shorter or somewhat longer primers, or primer sequences  
15 containing e.g. inositol residues or ambiguous bases or even primers that contain one or more mismatches when compared to the sequences of the primers of Figure 1. In general, sequences that exhibit at least 65%, more preferably at least 80% homology with the oligonucleotide sequences of Figure 1 are considered suitable for use in a method of the present invention.

20 Methods of the invention can in principle be performed by using any nucleic acid amplification method, such as the Polymerase Chain Reaction (PCR; Mullis 1987, U.S. Pat. No. 4,683,195, 4,683,202, en 4,800,159) or by using amplification reactions such as Ligase Chain Reaction (LCR; Barany 1991, Proc. Natl. Acad. Sci. USA 88:189-193; EP Appl. No., 320,308), Self-  
25 Sustained Sequence Replication (3SR; Guatelli *et al.*, 1990, Proc. Natl. Acad. Sci. USA 87:1874-1878), Strand Displacement Amplification (SDA; U.S. Pat. Nos. 5,270,184, en 5,455,166), Transcriptional Amplification System (TAS; Kwoh *et al.*, Proc. Natl. Acad. Sci. USA 86:1173-1177), Q-Beta Replicase (Lizardi *et al.*, 1988, Bio/Technology 6:1197), Rolling Circle Amplification  
30 (RCA; U.S. Pat. No. 5,871,921), Nucleic Acid Sequence Based Amplification

(NASBA), Cleavase Fragment Length Polymorphism (U.S. Pat. No. 5,719,028), Isothermal and Chimeric Primer-initiated Amplification of Nucleic Acid (ICAN), Ramification-extension Amplification Method (RAM; U.S. Pat. Nos. 5,719,028 and 5,942,391) or other suitable methods for amplification of DNA.

5 A preferred embodiment according to the present invention uses a PCR method comprising the primer-initiated amplification of supergroup B HPV-DNA with a set of 5 overlapping forward and 4 overlapping reverse primers that together match sequences within the L1 open reading frame of all known HPV types belonging to supergroup B.

10 Since the two consensus regions in the L1 ORF as defined herein are closely spaced together, a preferred set of bi-directional amplification primers amplifies a short fragment (i.e. about 72 base pairs) of supergroup B HPV-DNA. This provides for the possibility to amplify DNA sequences from template DNA that is of poor quality, such as DNA that is degraded to a  
15 certain extent, which allows for the use of formalin fixed specimens as a sample material, as well as of other less well preserved materials for use in a method of the present invention.

In order to amplify the DNA of HPV types with a small number of mismatches with one or more of the amplification primers, an amplification  
20 reaction may be performed under conditions of reduced stringency (e.g. a PCR amplification using an annealing temperature of 38°C, or the presence of 3.5 mM MgCl<sub>2</sub>). The person skilled in the art will be able to select conditions of suitable stringency.

Primer sequences and corresponding sequences of supergroup B HPV-  
25 specific DNA amplification primers are shown in Figure 1.

Another important aspect of the sequence region flanked by the two consensus regions in the L1 ORF is that the DNA sequence thereof comprises common motifs for various HPVs belonging to supergroup B, as can be seen in Figure 4.

Therefore, by amplifying the DNA sequence region between the two consensus regions in the L1 ORF that are specific for supergroup B HPVs, all supergroup B HPVs may be detected by using a set of generic cutaneous supergroup B HPV detection probes that essentially will hybridise to the amplicons of those HPVs that contain the common motif. When used in concert, such generic detection probes will bind all supergroup B HPVs amplification products. The inventors have found that a very advantageous system employs the hybridization to a cocktail of 3 overlapping generic probes provided with wobbles to detect all possible HPV types amplified by the cutaneous supergroup B HPV primers. An example of a set of suitable generic detection probes that binds to the region between the above mentioned two consensus regions of amplicons of all supergroup B HPVs is presented in Figure 3.

From the present disclosure it will be apparent to the person skilled in the art that other sets of generic detection probes than those of Figure 3 may be designed and used for detection of the DNA sequence flanked by the two consensus regions in the L1 ORF of all cutaneous supergroup B HPVs. For instance, it is possible to use somewhat shorter or somewhat longer probes, or probe sequences containing ambiguous bases or wobbles that are directed to a somewhat different cluster of HPV types, or an otherwise different set of probes than those of Figure 3, such as probes that contain one or more mismatches to their target sequence. In general, sequences that exhibit at least 65%, more preferably at least 80% homology with the oligonucleotide sequences of Figure 3, are considered suitable for use in a method of the present invention for the generic detection of amplicons of all supergroup B HPVs.

Also the complementary sequences of those listed in Figure 3 may suitably be used as generic detection probes in a method of the invention, provided that the complementary strand is amplified in the amplification reaction employed, such as when e.g. PCR is used for amplification.



The detection of the amplification products can in principle be accomplished by any suitable method known in the art. The detection fragments may be directly stained or labelled with radioactive labels, antibodies, luminescent dyes, fluorescent dyes, or enzyme reagents. Direct  
5 DNA stains include for example intercalating dyes such as acridine orange, ethidium bromide, ethidium monoazide or Hoechst dyes.

Alternatively, the DNA fragments may be detected by incorporation of labelled dNTP bases into the synthesized DNA fragments. Detection labels which may be associated with nucleotide bases include e.g. fluorescein, cyanine  
10 dye or BrdUrd.

The present invention preferably involves the detection of the reaction products obtained by the above described DNA amplification reaction on cutaneous supergroup B HPV DNA by hybridising the reaction products to one or more generic cutaneous HPV supergroup B probes.

15 When using a probe-based detection system, a suitable detection procedures for use in the present invention may for example comprise an enzyme immunoassay (EIA) format (Jacobs *et al.*, 1997, J. Clin. Microbiol. 35, 791-795). For performing a detection by manner of the EIA procedure, either the forward or the reverse primer used in the amplification reaction may  
20 comprise a capturing group, such as a biotin group for immobilization of supergroup B HPV-DNA PCR amplicons on e.g. a streptavidin coated microtiter plate wells for subsequent EIA detection of supergroup B HPV-amplicons (see below). The skilled person will understand that other groups for immobilization of supergroup B HPV-DNA PCR amplicons in an  
25 EIA format may be employed.

#### Typing of supergroup B HPVs

Another important aspect of the DNA sequence region flanked by the two consensus regions in the L1 ORF of the cutaneous supergroup B HPV is

that the DNA sequence thereof is also unique to each HPV type belonging to supergroup B, as can be seen in Figure 4.

Therefore, and in another aspect of the invention, the various HPV types belonging to supergroup B may be detected by using a plurality of type-specific cutaneous supergroup B HPV detection probes that, individually, bind  
5 only to the amplicon of one specific type of HPV belonging to supergroup B. Thus, by amplifying the DNA sequence region between the two consensus regions as defined herein, followed by probing for the unique sequences of each individual HPV-type, a method is provided for typing of the various  
10 supergroup B HPVs.

Probes useful for the type-specific detection of the region in the L1 ORF from DNA of supergroup B-type HPVs as disclosed herein are exemplified in Figure 5. These type-specific probes preferably bind only to at least a part of the DNA sequence region as amplified between the two consensus regions of a  
15 single HPV-type belonging to supergroup B. Those of skill in the art can prepare similar probes for type-specific detection based on the nucleotide sequence of the L1 ORF from the DNA of supergroup B HPVs without undue experimentation as set out herein. Also the complementary sequences of those listed in Figure 5 may suitably be used as type-specific detection probes in a  
20 method of the invention, provided that such a complementary strand is amplified in the amplification reaction employed.

Probes used in a method of typing according to the invention bind selectively to the amplicon of a certain HPV type or types, but not - or only weakly under hybridization conditions of very low stringency - to the amplicon  
25 of other HPV types. Accordingly, in one embodiment, the invention can be used to determine the type or types of HPV infecting a patient. This is very significant, as progression to malignant disease (and hence clinical prognosis) is heavily dependent on HPV type. Accordingly, in a typing aspect, the invention provides a method of determining the type(s) of HPV infection in a  
30 patient.

In a method of typing of supergroup B HPVs according to the invention, suitable probes are for instance those that hybridize to the amplicons of only a subset of HPV types. Such probes may for instance be the generic probes of Figure 3. In such an embodiment, for example, the ability to  
5 limit the infecting HPV type(s) to a particular subset (or exclude such a subset) may be sufficient.

In a method of typing of supergroup B HPVs according to the invention, the subset of amplicons to which a typing probe binds may consist of a single HPV type amplicon, or may consist of a plurality of amplicons of  
10 different types, such that binding or non-binding (as appropriate) of the type-specific probe to the amplicons generated from the sample will allow an investigator to make certain deductions about the identity of the HPV type(s) infecting the patient. Probes which bind to desired HPV types, but not to undesired HPV types, may be generated by methods known to the skilled  
15 person.

The present invention now provides a particular region of the amplicon to which probes may bind with considerable specificity. Although homologous regions may exist in all HPV amplicons generated, the nucleic acid sequence of this particular region is unique for HPVs of different type.

20 If a similar typing needs to be performed for supergroup B HPVs which are not (yet) included or discovered, clearly, the nucleic acid sequence of the amplicon of such HPV types will not necessarily be complementary to that of any of the probes herein provided. However, with the benefit of the present disclosure a suitable hybridization region can readily be identified in other  
25 amplicons by those skilled in the art by use of conventional alignment and sequence comparison computer programs.

In a preferred embodiment of a method for typing of supergroup B HPVs of the present invention a set of 24 different probes (Figure 5), or the complementary sequences thereof, is used that hybridise to the type-specific

sequences in the amplicons derivable from the 24 different cutaneous supergroup B HPVs as shown in Figure 4.

Suitable detection procedures for use in the typing aspect of present invention may for example comprise immobilization of the amplicons and  
5 probing the DNA sequences thereof by e.g. southern blotting. Other formats may comprise an EIA format as described above. To facilitate the detection of binding, the type-specific supergroup B HPV probes may comprise a label moiety such as a fluorophore, a chromophore, an enzyme or a radio-label, so as to facilitate monitoring of binding of the probes to the reaction product of the  
10 amplification reaction. Such labels are well-known to those skilled in the art and include, for example, fluorescein isothiocyanate (FITC),  $\beta$ -galactosidase, horseradish peroxidase, streptavidin, biotin, digoxigenin,  $^{35}\text{S}$  or  $^{125}\text{I}$ . Other examples will be apparent to those skilled in the art.

Although other methods may be employed, a method of typing  
15 according to the invention is preferably performed by a so called reverse line blot (RLB) assay, such as recently developed for e.g. mucosotropic HPVs (van den Brule *et al.*, 2002, J. Clin. Microbiol. 40, 779-787). For this purpose RLB probes are preferably synthesized with a 5' amino group for subsequent immobilization on e.g. carboxyl-coated nylon membranes. The advantage of an  
20 RLB format is the ease of the system and its speed, thus allowing for high throughput sample processing.

A method of typing of supergroup B HPVs according to the invention may be performed on similar sample material as that used for the detection of supergroup B HPVs described above. Alternatively, the typing of the  
25 supergroup B HPV according to a method of the invention may be performed directly on amplicons or DNA amplification reaction products obtained by using the DNA amplification reaction as described for the detection of supergroup B HPV herein above.

As the difference between the two aspects of a method of the  
30 invention is in the application of different detection probes, a method of the

invention may in one aspect comprise the amplification of suspected supergroup B HPV DNA, followed by the detection of amplicons of any HPV (i.e. of an infection with supergroup B HPV) by using a generic detection probe cocktail that, in concert, hybridises to all supergroup B HPV amplicons generated. In another aspect, a method of the invention may comprise the amplification of supergroup B HPV DNA, followed by the typing for 24 HPVs in, for example a reverse line blot assay. Thus, detection and typing according to methods of the invention may be performed in parallel.

#### 10 Primers and probes

In a further aspect, the invention provides a set of bi-directional amplification primers for the amplification of supergroup B HPV-DNA for use in a method of the invention. These primers are preferably oligonucleotide primers with a sequence length of about 8 to about 35, preferably of about 15 to about 30, more preferably of about 21 to about 28 nucleotides. In a most preferred embodiment the forward primer comprises about 21 nucleotides, while the reverse primer comprises about 28 nucleotides. Preferably one of the primers is labelled with a group to facilitate the immobilization of one of strands of the double stranded supergroup B HPV amplicons and in an even more preferred embodiment the reverse primer is labelled with biotin.

The nucleotide sequence of the bi-directional primers of the present invention should be designed to allow the formation of a stable hybrid between the primer and the target DNA at the position of one of the two consensus regions in the L1 ORF of the supergroup B HPV genome as defined herein. Thereto, the nucleotide sequence of the forward primer is preferably substantially complementary to the anti-sense strand of the L1 ORF in the genome of one or more supergroup B HPVs in the region defined as forward primer region in Figure 2, while the nucleotide sequence of the reverse primer is preferably substantially complementary to the sense strand of the L1 ORF in the genome of one or more supergroup B HPVs in the region defined as

reverse primer region in Figure 2. Primers that are consistent with this criterion are for example those displayed in Figure 1. When used in concert, the primers of Figure 1 will amplify DNA of a corresponding region in the genome of all supergroup B HPVs.

5           The primers herein are selected to be "substantially" complementary (i.e. at least 65%, more preferably at least 80% perfectly complementary) to the consensus regions present on the different strands of each specific sequence to be amplified. It is possible to use somewhat shorter or somewhat longer primers, or primer sequences containing e.g. inositol residues or ambiguous  
10   bases or even primers that contain one or more mismatches when compared to the sequences of the primers of Figure 1. In general, sequences that exhibit at least 65%, more preferably at least 80% homology with the oligonucleotide sequences of Figure 1, are considered suitable for use in a method of the present invention. Sequence mismatches are also not critical when using low  
15   stringency hybridization conditions. In the present invention, the primers are preferably substantially complementary to the sequence of at least one of the two consensus regions in the L1 ORF of supergroup B HPVs.

          In another aspect, the invention provides oligonucleotide probes for the generic detection of supergroup B HPV-DNA. The detection probes herein  
20   are selected to be "substantially" complementary to one of the strands of the double stranded DNA amplicons generated by an amplification reaction of the invention. Preferably the probes are substantially complementary to the immobilizable, e.g. biotin labelled, antisense strands of the amplicons generated from the L1 ORF of the supergroup B HPVs.

25           The amplicons comprise regions characterized by common motifs as well as unique sequences between the various supergroup B HPVs. The generic probes for detection of these amplicons are designed to be sufficiently complementary to hybridise with either strand of the amplicons generated from one or more supergroup B HPV types. The generic detection probe  
30   sequence need not reflect the exact sequence of either strand of the amplicons,

nor does the generic detection probe sequence need to reflect the exact sequence of all supergroup B HPVs. The present invention now provides in Figure 3 a set of probes with ambiguous bases (wobbles) to facilitate their binding to at least a subset of all possible amplicons generated from the various supergroup B HPVs. This minimizes the number of generic probes required for detection of all supergroup B HPVs.

It is allowable for generic detection probes of the present invention to contain one or more mismatches to their target sequence. In general, sequences that exhibit at least 65%, more preferably at least 80% homology with the oligonucleotide sequences of Figure 3, are considered suitable for use in a method of the present invention for the generic detection of amplicons of all supergroup B HPVs. Therefore, when referring to the primers or probes of Figures 1 and 3, it is intended that sequences exhibiting at least 65%, more preferably at least 80% homology with those of Figures 1 and 3 are included.

Also the complementary sequences of those listed in Figure 3 may be used as generic detection probes in a method of the invention, provided that the complementary strand is amplified in the amplification reaction employed.

Probes which bind to some supergroup B HPV types, but not to other supergroup B HPV types, may be generated by methods known to the skilled person.

To facilitate detection the generic supergroup B HPV probes may comprise a label moiety such as a fluorophore, a chromophore, an enzyme or a radio-label, so as to facilitate monitoring of binding of the probes to the reaction product of the amplification reaction. Such labels are well-known to those skilled in the art and include, for example, fluorescein, cyanine dye,  $\beta$ -galactosidase, horseradish peroxidase, streptavidin, biotin,  $^{35}\text{S}$  or  $^{125}\text{I}$ . Other examples will be apparent to those skilled in the art. A preferred label for a generic supergroup B HPV probe used in a method of the present invention is a digoxigenin (DIG) label.

These generic detection probes are preferably oligonucleotide probes with a sequence length of about 8 to about 70, preferably of about 15 to about 40, more preferably of about 25 to about 35 nucleotides. In a most preferred embodiment the generic detection probes comprise about 30 nucleotides.

5           In yet another aspect, the invention provides oligonucleotide probes for the type-specific or selective detection of individual HPV-types or subsets of types belonging to supergroup B of HPV. In a preferred embodiment one probe is provided for each HPV type, such probes therefore being type-specific. The present invention discloses a set of 24 suitable type-specific probe sequences  
10 for the type-specific detection of the specific supergroup B HPVs mentioned in Figure 5. Although some mismatches would be allowable it is preferred that the typing probes are perfectly complementary to their respective target sequences. As outlined above, the complementary sequences of those presented in Figure 5 may also be used as type specific probes in methods of the present  
15 invention, provided that the amplification reaction allows for the amplification of the complementary DNA fragments. Therefore, when referring to the probes of figures 3 and 5, it is intended that complementary sequences thereof are included.

          Type-specific or selective detection probes can be used in a variety of  
20 typing procedures. Suitable procedures comprise those whereby the probe-target is immobilized, but also procedures whereby the probes are immobilized may be used. As examples of the latter, both biochip-type of formats, but also reverse line blot formats, such as RLB or Line Probe Assays may be used. Depending on the substrate for immobilization and the surface chemistry  
25 selected, the skilled person is able to chose a suitable group with which the oligonucleotide probe is labelled for substrate immobilization. In a preferred embodiment this group is an amino-group introduced at the 5'-end of the probe.

          The type-specific detection probes are preferably oligonucleotide  
30 probes with a sequence length of about 8 to about 30, preferably of about 15 to



about 25, more preferably of about 19 to about 23 nucleotides. As with the design of the generic detection probes, the skilled person is aware that the annealing temperature of individual probes within a set of probes that is used in one hybridization reaction may suitably be adjusted by providing balance  
5 between G+C content and length of the probe.

In one embodiment the present invention also provides a kit comprising primers and probes for performing a method of the invention. A preferred embodiment of such a kit comprises both the amplification primers, the generic detection probes and the type-specific probes as disclosed in  
10 Figures 1, 3, and 5, respectively.

The following non-limiting examples illustrate several embodiments of the present invention.

#### Example 1.

15 Amplification of HPVs of supergroup B, EIA detection and RLB typing.

#### *Methodology*

##### HPV test panel.

For the evaluation of the sensitivity and specificity of the RLB typing  
20 method, plasmid clones of the following HPV types were used as test panel:  
HPV 4, 5, 8, 9, 14, 17, 19, 21, 22, 23, 25, 36, 38, 49, and 50.

##### Clinical specimens.

For the evaluation of HPV detection and typing on clinical specimens, formalin-fixed, paraffin embedded tissues of skin samples were used. A series  
25 of 5 µm sections were cut using the so-called sandwich method (Walboomers et al., 1999, J. Pathol. 189:12-19). Briefly, outer sections were haematoxylin-eosin stained for histological analyses, whereas inner sections (in total approximately 1 cm<sup>2</sup> of tissue) were placed into a tube and used for PCR purposes. 250 µl lysis mix (10 mM Tris-HCl (pH 7.5) , 0.45% Tween 20 and 500  
30 µg proteinase K (Roche, Mannheim, Germany) ) was added and incubation was

performed overnight at 37°C. The sample was boiled and centrifuged 1 min in a standard table top centrifuge (10,000 rpm). After cooling 10 µl was used for PCR.

#### Cutaneous HPV supergroup B-PCR.

5            Isolated HPV plasmid DNA or pre-treated samples were subjected to PCR using a mixture of 5 forward primers (i.e. 5f, 6f, 7f, 8f, 9f; Figure 1) and 3 biotinylated reverse primers (i.e. 5r-bio, 6r-bio, 7r-bio, and 8r-bio; Figure 1). The PCR reaction mixtures of 50 µl contained 1.5 mM MgCl<sub>2</sub>, 200 µM of each dNTP, 12.5 pmol of each primer, 1 unit AmpliTaq Gold DNA polymerase  
10            (Perkin-Elmer, Foster City, CA), and various concentrations of HPV plasmid DNA or 10 µl of crude sample extract, in Taq Gold buffer (Perkin-Elmer, Foster City, CA). 40 PCR cycles were performed using a PE 9700 thermocycler (Perkin-Elmer, Foster City, CA). Each PCR reaction was initiated by a preheating step for 10 minutes at 94°C followed by 40 cycles consisting of a  
15            denaturation step (20 seconds at 94°C), a primer annealing step (30 seconds at 38°C), and an elongation step (80 seconds at 71°C). Ramping times were as described before (van den Brule et al., 2002, J. Clin. Microbiol. 40: 779-787). The final elongation step was extended for 4 minutes.

#### Enzyme-Immunoassay of PCR products.

20            The EIA was performed using streptavidine-coated microtitre plates (Roche, Mannheim, Germany) and a read-out after overnight incubation as previously described (Jacobs et al., 1997, J. Clin. Microbiol. 35:791-795), except that a hybridisation temperature of 55°C was used. A mixture of 3 DIG-labelled generic oligoprobes (ie. SkinPR 1-DIG, SkinPR 2-DIG, and SkinPR3-  
25            DIG; Figure 3) was used for hybridisation.

#### HPV reverse line blotting (RLB) of PCR products.

RLB analyses were performed using a system that previously has been developed for mucosotropic HPV types (van den Brule et al., 2002, J. Clin. Microbiol. 40: 779-787). The system is based on the use of a miniblottedter for  
30            spotting in parallel up to 42 different oligoprobes containing a 5'-amino group

on a carboxyl-coated nylon membrane. Subsequently, up to 42 PCR products can be pipetted into the parallel channels of the miniblotted in such way that the channels are perpendicular to the rows of oligoprobes deposited previously. In this format hybridisation is carried out, followed by incubation of the  
5 membrane with anti-biotin conjugate and Enhanced Chemiluminescence (ECL) detection.

The HPV-specific 5' amino-linked oligonucleotides (Isogen, Maarsse, The Netherlands) selected for RLB are shown in Figure 5. Selection was performed in such a way that  $T_m$  was (approximately) 55°C and the length was  
10 approximately 20 bp using primer/probe selection software (Oligo). The oligoprobes were covalently bound to negatively charged membranes (Biodyne C, Pall BioSupport, East Hills, NY). Briefly, membranes were activated by using 16 % (wt/vol) EDAC (1-ethyl-3-(3-dimethylaminopropyl)carbodiimide; Sigma, St. Louis, MO) and placed on a miniblotted system (MN 45,  
15 Immunitics, Cambridge, MA). The oligonucleotides were applied in parallel lines and after a 1 minute incubation the channels were aspirated and the membrane inactivated using 100 mM NaOH followed by washing in 2x SSPE (0.18 M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub> and 1 mM EDTA (pH 7.7); Life Technologies, Rockville, MD) supplemented with 0.1% SDS (BDH, Dorset, England).

20 For hybridisation, 10 µl of biotinylated PCR product was diluted in 150 µl of 2x SSPE/0.1%SDS, heat-denatured (96°C) and rapidly cooled on ice. The PCR products were pipetted into the parallel channels perpendicular to the rows of oligoprobes. Hybridisation was performed at 42°C for 1 hour, followed by two washings in 2x SSPE/0.5%SDS at 51°C. Subsequently,  
25 membranes were incubated with streptavidin-peroxidase conjugate for 45 to 60 min at 42° C (1:4000 times diluted in 2x SSPE/0.5% SDS; Roche, Mannheim, Germany), washed twice in 2x SSPE/0.5%SDS for 10 min at 42°C, and rinsed twice in 2x SSPE for 5 min at room temperature. Detection was done, by using chemoluminescent ECL detection liquid (Amersham, Buckinghamshire,  
30 England) followed by an exposure to a film (Hyperfilm, Amersham,

Buckinghamshire, England) for 1 to 10 min. Films were subsequently developed. For repeated use, the membranes were stripped by washing two times in 1% SDS at 80°C for 30 min followed by an incubation in 20 mM EDTA (pH 8) at room temperature for 15 min. Membranes were sealed in plastic and  
5 stored at 4°C until further use.

### *Results*

The amplification of HPVs of supergroup B and subsequent EIA detection of PCR products derived from 1 ng of cloned HPVs 4, 5, 8, 9, 14, 17,  
10 19, 21, 22, 23, 25, 36, 38, 49, and 50 revealed clear EIA signals in all cases, whereas no cross-reactivity was observed with human DNA. This indicates a high specificity. Using dilution lines of cloned HPVs 5, 8, 21, and 50 mixed with human placental DNA the cutaneous HPV supergroup B -PCR EIA also performed very well in terms of sensitivity; the analytical sensitivity ranged  
15 from 10 fg (HPV 5, and 21) to 100 fg (HPV 8 and 50). Also RLB typing revealed specific signals without cross-reactivity and the sensitivity was equal to if not higher than that of the EIA.

### Example 2.

#### 20 Analysis of formalin-fixed clinical specimens

The amplification of cutaneous supergroup B HPV DNA was tested using the materials and methods described in Example 1 except that DNA was used that was isolated from formalin-fixed and paraffin-embedded tissue samples from 87 cutaneous specimens that showed good DNA quality (i.e.  
25 positive samples in a beta-globin PCR). These samples included 59 samples of renal transplant recipients, comprising 21 cases of squamous cell carcinoma (SCC), 23 cases of basal cell carcinoma (BCC) and 15 cases of actinic keratosis (AK). Another 29 samples were obtained from immunocompetent patients, comprising 13 cases of SCC, 12 cases of BCC, and 4 cases of AK.

EIA detection with generic supergroup B probes revealed positivity in 62% of the cases: 76% of specimens from renal transplant recipients and 34% of specimens of immunocompetent patients. These included 15 out of 21 cases of SCC (71 %), 15 out of 23 cases of BCC (65%), and 14 out of 15 cases of AK (93%) in samples obtained from renal transplant recipients and 4 out of 13 cases of SCC (31 %), 2 out of 12 cases of BCC (17%) and 4 out of 4 cases of AK (100 %) in samples obtained from immunocompetent patients.

Typing by RLB of the SCC samples of both renal transplant recipients and immunocompetent patients revealed in between 1 and 7 different types of HPV per sample. All together, the following 19 HPV types were detected in one or more samples: HPV 4, 5, 8, 9, 14, 15, 17, 19, 20, 21, 23, 24, 25, 36, 37, 38, 47, 49, 50. These data are in agreement with data obtained from frozen specimens published in literature using other PCR systems (Shamani *et al.*, 1996, J. Natl. Cancer Inst. 88: 802-811; Berkhout *et al.*, J. Clin. Microbiol. 33: 690-695; Forslund *et al.*, J. Gen. Virol. 80: 2437-2443; Harwood *et al.*, J. Med. Virol. 61: 289-297; Meijer *et al.*, Cancer Detect. Prev. 25: 533-547).

Claims

1. Method for the detection of cutaneous supergroup B HPVs comprising the steps of:
  - (a) providing a sample suspected of harbouring cutaneous supergroup B HPVs;
  - 5 (b) providing a plurality of pairs of bi-directional primers collectively substantially complementary to DNA of all cutaneous supergroup B HPVs;
  - (c) performing a reaction to amplify DNA derived from the said sample using said plurality of primers; and
  - 10 (d) detecting DNA amplification products from cutaneous supergroup B HPV from said sample.
2. Method according to claim 1, wherein step (d) is carried out by hybridising the reaction products of the said DNA amplification reaction to a plurality of generic cutaneous supergroup B HPV probes.
- 15 3. Method according to claim 1 or 2, wherein said plurality of pairs of bi-directional primers comprises primers that are collectively substantially complementary to a first consensus region in the DNA of all cutaneous supergroup B HPVs, and which plurality further comprises primers that are collectively substantially complementary to a second consensus region in the  
20 DNA of all cutaneous supergroup B HPVs.
4. Method according to claim 3, wherein said first and second consensus regions are in the L1 ORF of cutaneous supergroup B HPVs.
5. Method according to claim 4, wherein said first and second consensus regions are substantially as defined in Figure 2.
- 25 6. Method according to claim 5, wherein said plurality of pairs of primers comprises the primers of Figure 1.

7. Method according to claim 6, wherein one of each pair of primers of said plurality of pairs of bi-directional primers comprise a biotin label.
8. Method according to any one of the previous claims, wherein said reaction to amplify DNA is performed under conditions of reduced stringency.
- 5 9. Method according to any one of the claims 2-8, wherein said plurality of supergroup B HPV probes is substantially complementary to the nucleic acid sequence of the said DNA amplification products from all supergroup B HPVs.
- 10 10. Method according to any one of the claims 2-9, wherein said supergroup B HPV probes comprise the probes of Figure 3.
11. Method according to claim 10, wherein said probes comprise a DIG label.
12. Method for typing of cutaneous supergroup B HPVs comprising the steps of:
  - 15 (a) providing DNA amplification products by amplifying DNA of cutaneous supergroup B HPV using a plurality of pairs of bi-directional primers; and
  - (b) detecting DNA amplification products from one or more supergroup B HPV types by hybridising the said amplification products to at least one cutaneous supergroup B HPV probe that  
20 is substantially complementary to the DNA of at least one but not all cutaneous supergroup B HPV types.
13. Method according to claim 12, wherein said plurality of pairs of bi-directional primers comprises primers that are collectively substantially  
25 complementary to a first consensus region in the DNA of all cutaneous supergroup B HPVs, and which plurality further comprises primers that are collectively substantially complementary to a second consensus region in the DNA of all cutaneous supergroup B HPVs.
14. Method according to claim 13, wherein said first and second  
30 consensus regions are in the L1 ORF of cutaneous supergroup B HPVs.

15. Method according to claim 14, wherein said first and second consensus regions are substantially as defined in Figure 2.
16. Method according to claim 15, wherein said plurality of pairs of bi-directional primers comprises the primers of Figure 1.
- 5 17. Method according to claim 16, wherein one of each pair of primers of said plurality of pairs of bi-directional primers comprise a biotin label.
18. Method according to any one of claims 12-17, wherein said at least one probe is substantially complementary to the DNA of exactly one type of supergroup B HPV.
- 10 19. Method according to any one of the claims 12-18, wherein said at least one probe is selected from the probes of Figure 5.
20. Method according to any one of the claims 12-19, wherein the detection comprises the use of a reverse line blot.
21. Bi-directional primers for use in a method according to any one of
- 15 the previous claims, which primers are collectively substantially complementary to a first and a second consensus region in the L1 ORF of all supergroup B HPVs.
22. Bi-directional primers of Figure 1.
23. Generic detection probes for the detection of cutaneous supergroup B
- 20 HPVs, which probes are collectively substantially complementary to a region in the L1 ORF of all supergroup B HPVs between nucleotide positions 6539 and 6610 of HPV 4 and corresponding region of the other cutaneous supergroup B HPV types.
24. Generic detection probes of Figure 3.
- 25 25. Detection probes for the detection of cutaneous supergroup B HPV types, which probes are substantially complementary to a region in the L1 ORF of at least one but not all cutaneous supergroup B HPV types between nucleotide positions 6539 and 6610 of HPV 4 and corresponding region of the other cutaneous supergroup B HPV types.



26. Type-specific detection probes for the detection of cutaneous supergroup B HPV types, which probes are substantially complementary to a region in the L1 ORF of exactly one type of cutaneous supergroup B HPV type between nucleotide positions 6539 and 6610 of HPV 4 and corresponding region of the other cutaneous supergroup B HPV types.
27. Type-specific detection probes of Figure 5.

Title: Method for detecting and typing of cutaneous HPV and primers and probes for use therein.

Abstract

The present invention relates to a method for detecting and typing of human papillomavirus (HPV). In particular, the invention relates to a method for detecting and typing cutaneous HPVs of supergroup B and to primers and probes for use in a method of the invention. An important advantage of the present invention is that a large number of HPV types can be detected in a single assay, that it is applicable to fixed and poorly preserved material and that the detection method and typing method can be coupled.

**Forward primers (5'- 3')**

5f: TTAGCTCAAATTAATGCAATG  
6f: TTGGCACAAATTAATGCAATG  
7f: TTAGCACAAATTAATGCTATG  
8f: TTAGCTCAGATAAATGCAATG  
9f: CTGGCTCATTAAATGTAATG

**Reverse primers (5'bio-3')**

5r-bio: ATCAGGGGTGGGTACAAATCCTAACTGC  
6r-bio: ATCAGGGGTGGGTACAAATCCTAATTGC  
7r-bio: ATCTGGTGCAGGAACAAATCCTAATTGC  
8r-bio: GGGAGGTGGTGGAAACAAAAGCTAATTGC

**Figure 1**

Type	forward primer region (sense)	reverse primer region (anti-sense)
HPV 4:	ATGGCTCATTTAAATGTAATG	GGGAGGTGGTGGAAACAAAATTTAACTGC
HPV 5:	TTGGCACAGATCAATGCAATG	ATCAGGAGTGGGAACAAATCCTAACTGC
HPV 8:	TTAGCACAAATCAATGCCATG	ATCAGGGGTAGGTACAAATCCCAGTTGC
HPV 9:	TTATCCCAGATAAATGCAATG	TTCAGGTGATGGAACAAACCCTAATTGC
HPV 12:	TTAGCACAGATAAATGCAATG	GTCAGGAGTTGGCACGAAGCCTAATTGC
HPV 14:	TTAGCTCAAATTAATGCAATG	GTCTGGTGCAGGTACAAATCCTAATTGC
HPV 15:	TTAACACAAATTAATGCTATG	GTCTGGTGTAGGAACAAACCCTAGTTGC
HPV 17:	TTAACTCAAATTAATGCAATG	ATCTGGTGTAGGCACAAATCCTAACTGC
HPV 19:	CTGGCACAAATCAATGCAATG	ATCGGGTGCAGGCACGAAACCTAATTGC
HPV 20:	TTAGCTCAAATTAATGCTATG	ATCCGGTGCAGGAACGAATCCTAACTGC
HPV 21:	TTAGCTCAAATTAATGCAATG	GTCTGGGGCAGGAACAAATCCTAACTGC
HPV 22:	TTAACTCAAATTAATGCCATG	GTCTGGTGTAGGTACAAAGCCTAGTTGC
HPV 23:	TTAACACAAATTAATGCCATG	ATCTGGTGTAGGAACAAACCCTAACTGC
HPV 24:	TTAGCACAGATCAATGCTATG	GTCTGGTGCAGGCACAAATCCTAACTGC
HPV 25:	TTGGCTCAGATTAATGCAATG	GTCCGGTGCAGGAACGAATCCTAATTGC
HPV 36:	TTGGCTCAGATAAATGCTATG	ATCTGGAGTAGGTACAAAACCTAACTGC
HPV 37:	TTAACTCAGATAAATGCAATG	ATCTGGAGTAGGTACAAATCCTAATTGC
HPV 38:	CTGACACAGATTAATGCTATG	GTCTGGGTGGGGTACAAAGCCTAATTGC
HPV 47:	TTAGCACAAATTAATGCCATG	GTCTGGAGTAGGCACAAATCCTAACTGC
HPV 48:	CTGGCACATTTACACGCTTTG	TGGAGCTGGTGGAAACAAAAGCTAATTGC
HPV 49:	CTGGCACAAATCAATGCTATG	ATCAGGGGTAGGAACAAATCCCAATTGC
HPV 50:	TTAGCTCATCTTAATGTAATG	GGGAGCTGGTGGAACTAAAGCTAATTGC
HPV 60:	TTAGCTCATTTAAATGTAATG	TGGTGGTGGAGGTACAAATGAAAGTTGC
HPV 65:	ATGGCTCACTTAAATGTAATG	AGGAGGAGGGGGCACAAAATTTAATTGC

Figure 2

SkinPR 1-DIG : GCAATGAATSCHAAAYATWTTRGARGADTGG  
SkinPR 2-DIG : GCAATGAAYTCWDVTATRTTGGARGATTGG  
SkinPR 3-DIG : GCAATGAAYYCHDVDHTWTTRGARGADTGG

D = A, T or G  
H = A, T or C  
R = A or G  
S = C or G  
V = A, C or G  
W = A or T  
Y = C or T

**Figure 3**

HPV 4 : GTAATGAATCCTAATATTTTGGATAATTGG  
HPV 5 : GCAATGAACTCTTCGTTATTGGAGGATTGG  
HPV 8 : GCCATGAATTCATCACTACTTGAGGAATGG  
HPV 9 : GCAATGAACTCAGGTATTTTAGAGGATTGG  
HPV 12 : GCAATGAACTCTTCTTTGCTGGAGGACTGG  
HPV 14 : GCAATGAATTCTAATATTTTAGAGGAGTGG  
HPV 15 : GCTATGAATTCAGGTATTTTAGAAGACTGG  
HPV 17 : GCAATGAACTCAGGAATTCTGGAAGACTGG  
HPV 19 : GCAATGAATTCTAACATATTGGAAGAATGG  
HPV 20 : GCTATGAATTCAAATATATTAGAGGAGTGG  
HPV 21 : GCAATGAATGCAAATATTTTAGAAGAATGG  
HPV 22 : GCCATGAATCATGGCATTTTAGAAAATTGG  
HPV 23 : GCCATGAATTCAGATATTTTAGAGAATTGG  
HPV 24 : GCTATGAATCCATCTCTATTGGAAGAGTGG  
HPV 25 : GCAATGAATTCCAACATTTTAGAAGATGG  
HPV 36 : GCTATGAATTCTTCTTTATTGGAAGATTGG  
HPV 37 : GCAATGAATTCTGGTATATTGGAAGAGTGG  
HPV 38 : GCTATGAATTCTGGAATATTAGAAAATTGG  
HPV 47 : GCCATGAATTCGTCTCTTTTAGAGGAATGG  
HPV 48 : GTTATGAATCCCAGAATATTAGAGGAGTGG  
HPV 49 : GCTATGAATTCTTCTATATTGGAAAATTGG  
HPV 50 : GTAATGGACCCCAGAATTTTGGAAAATTGG  
HPV 60 : GTAATGGACAAAAATATTTTAGAGGATTGG  
HPV 65 : GTAATGAATCCTAATATCTTAGACAATTGG

**Figure 4**

RLBskinHPV 4 : AATCCTAATATTTTGGATAATTG  
RLBskinHPV 5 : TCTTCGTTATTGGAGGATTG  
RLBskinHPV 8 : AATTCATCACTACTTGAGGA  
RLBskinHPV 9 : TCAGGTATTTTAGAGGATTG  
RLBskinHPV 12 : TTCTTTGCTGGAGGACTG  
RLBskinHPV 14 : ATTCTAATATTTTAGAGGAGTG  
RLBskinHPV 15 : AATTCAGGTATTTTAGAAGACT  
RLBskinHPV 17 : AACTCAGGAATTCTGGAAGA  
RLBskinHPV 19 : TTCTAACATATTGGAAGAATG  
RLBskinHPV 20 : ATTCAAATATATTAGAGGAGTG  
RLBskinHPV 21 : ATGCAAATATTTTAGAAGAATG  
RLBskinHPV 22 : AATCATGGCATTTTAGAAAATT  
RLBskinHPV 23 : ATTCAGATATTTTAGAGAATTG  
RLBskinHPV 24 : CATCTCTATTGGAAGAGTG  
RLBskinHPV 25 : AATTCCAACATTTTAGAAGAAT  
RLBskinHPV 36 : TTCTTCTTTATTGGAAGATTG  
RLBskinHPV 37 : TTCTGGTATATTGGAAGAGT  
RLBskinHPV 38 : ATTCTGGAATATTAGAAAATTG  
RLBskinHPV 47 : AATTCGTCTCTTTTAGAGGA  
RLBskinHPV 48 : CCCAGAATATTAGAGGAGT  
RLBskinHPV 49 : ATTCTTCTATATTGGAAAATTG  
RLBskinHPV 50 : ACCCCAGAATTTTGGAAAAT  
RLBskinHPV 60 : GACAAAAATATTTTAGAGGATT  
RLBskinHPV 65 : AATCCTAATATCTTAGACAATT

**Figure 5**




Figure 6

